

Extraction of Essential Oils from Afghanistan Medicinal Plants Using Microwave and Conventional Methods

Nazifa Faqeryar and Yoshihito Mori

Department of Chemistry and Biochemistry, Division of Advanced Sciences, Graduate School of Humanities and Sciences, Ochanomizu University 2-1-1 Otsuka, Bunkyo, Tokyo 112-8610, Japan

Abstract: Afghanistan has many kinds of volatile-oil-rich and not-fully-utilized flora. This article represents a review on Afghanistan medicinal plants contain essential oils. Lamiaceae, Asteraceae and Apiaceae are the most important botanical families which are widely found and used as traditional medicines, food and spices in Afghanistan. Over 215 plants which are used in traditional therapy in Afghanistan, 93 plants contain essential oils. Twenty-two plants contain essential oils applied to microwave extraction techniques and forty-nine plants contain essential oils applied to conventional extraction techniques. Totally 41 extracted species belong to above mentioned botanical families that could be found in Afghanistan too. But still there are some species of these families which are endemic in Afghanistan and not studied yet.

Key words: Afghanistan medicinal plants, essential oils, conventional extraction, microwave extraction.

1. Introduction

Afghanistan is a mountainous country which has a dry climate with very hot summers and very cold winters (+51 and -52.2 °C). This fluctuation of climate has effect on habitude of exploitation of agricultural land and the usage cycle of plants. Afghanistan has more than 5,000 plant species. About 1,200 plant species are endemic [1]. And a great number of plant species are used in traditional medicines [2]. Also most of these floras are volatile-oil-rich but not-fully-utilized.

Traditional medicines are the oldest medical practices in societies which are used long before modern medicines. Different nations, cultures and custom of different nations beholden their growth and development to the use of medicinal plants. According to WHO (World Health Organization) more than 80% of world's population used medicinal plants. It is not just in developing countries but also in USA, UK and other developed countries the usage of herbal

medicines is growing up.

In Afghanistan also traditional medicinal therapy has been used since centuries. People of Wakhan Corridor, in Pamir mountain of Afghanistan one of the most remote and insular areas in the world is using local herbal medicines for treatment of infectious disease, fever and pain [3]. Different way of usage of medicinal plants is common in Afghanistan. The most common traditionally way is boiling herbs or making herbal tea by decoction of flowers, leaves or stems of plant in water and then the extract is filtered, while numerous people are in unscientifically manner treat their health problems with plants. Villagers in Pamir mountain of Afghanistan and Tajikistan are using fresh and dried medical plants for treatment and prevention of gastrointestinal, dermatological, kidney illness and hypotension, pain relief etc. [4].

The aim of this study is to show the importance of Afghanistan medicinal plants and its usage for both domestic and industrial people in future. In this article we have reviewed two methods of extraction: microwave and conventional applied widely for extraction of essential oil from plant. Furthermore, the

Corresponding author: Nazifa Faqeryar, Ph.D., assistant professor, research fields: botanical chemistry and pharmaceutical chemistry.

spread of essential oil in three most important botanical families in the world (Lamiaceae, Apiaceae and Compositae), as well as the native species of Afghanistan belong to some great genus of these families and contain essential oil is discussed here.

2. Essential Oil

Essential oils in plants are complex mixtures of volatile substances, presented at low concentrations and broadly used in folk medicine for treatment and debarment of different human illness, as well as in the perfume industries especially in production of shampoos, hair lotions, bathing lotion and as disinfectants also in pharmaceutical sectors and in the food and human nutrition [5-7].

The yield and chemical composition of essential oils of a single species of the genus are affected by following factors: Method of extraction, geography or edaphic of growth of plant, collecting period and environmental conditions. *Ocimum sanctum* shows an increase in essential oil in plant collected from region of Kerala compared two plants collected from Maharashtra of India. Also the amount of essential oil was high in sample collected in winter compared to sample collected in summer [8].

Essential oils have been extracted by hydro-distillation, steam distillation and soxhlet extraction from decades. These techniques take from hours to days and require plenty amount of water, therefore, resulting in the losses of volatile compounds, degradation of unsaturated or ester compounds through thermal or hydrolytic effects and toxic solvents residue in the extract. These drawbacks have led to the developing of new alternative techniques in extraction such as supercritical carbon dioxide, supercritical fluids, ultrasound and microwave extractions [9-11].

Microwaves are form of electromagnetic energy at frequency between 300 MHz and 300 GHz [12]. The frequency of 2.45 GHz is the most reputed and has a substantial effect on the rate of various chemical and food industries processes [13]. Advancements in

microwave extraction have resulted in new techniques for instance compressed air microwave distillation (CAMD), vacuum microwave hydro distillation (VMHD), microwave hydro-distillation (MWHd), solvent free microwave extraction (SFME), microwave-accelerated steam distillation (MASD), microwave steam distillation (MSD) and microwave hydro diffusion and gravity (MHD). Nowadays, more than 500 articles have been published on the topics of microwave extraction [14]. Utilization of microwaves in extraction process has demonstrated to diminish the extraction time and volume of solvent required. Therefore, these methods are environmentally friendly and consuming less energy compared to conventional methods [15].

3. Spread of Essential Oil in Botanical Families

Lamiaceae (Labiatae), Apiaceae (Umbelliferae) and Asteraceae (Compositae) are the most interesting botanical families and there are large numbers of publications which have been reported worldwide.

Essential oils occurred widely in the plant kingdom. In 64 plants families, about 400 species produced essential oil. From viewpoint of essential oil production, the most important families are Asteraceae (Compositae), Lamiaceae (Labiatae) and Apiaceae (Umbelliferae). Fifteen species of each above-mentioned family produce essential oils in a large scale. Fabaceae, Rutaceae, Lauraceae, Cupressaceae, Pinaceae, Zingiberaceae, Myrtaceae, Rubiaceae and Burseraceae are the other important families producing essential oils [16]. Same species of these families are found in Afghanistan too and are utilized in folk medicines largely.

Brackle et al. [1] have shown distribution of these botanical families in Afghanistan. Jeppesen and his co-workers have studied the antibacterial effect of some species of Lamiaceae and Compositae families from Pamir mountain of Afghanistan [3]. A regional research in Behsood in a remote and impoverished

district of Maidan-wardak province of Afghanistan showed that most people in this area have used plants for treatment of different disease which belonged to Lamiaceae, Apiaceae and Asteraceae families.

Seventy-seven botanical families are represented as medicinal plants of Afghanistan traditionally used by people [2]. Over them, thirty-seven families contain essential oil. Tables 1 and 2 show Afghanistan medical plants contain essential oil.

3.1 Lamiaceae (Labiatae)

Lamiaceae botanical family contains 210 genera and 3,500 species [1]. The Lamiaceae family is rich in aromatic species. Therefore, these plants are used as culinary herb, folk medicines and perfumes in many countries. In Lamiaceae family the volatile oil is distributed on the aerial parts of plant [17].

In Afghanistan, 234 species of Lamiaceae family are found in which 67 of them are endemic [1]. Fifteen plants of Lamiaceae family are introduced as medicinal plant [2]. Thirteen plants of this family contain essential oil. *Nepeta laevigata*, *Salvia hydrangea*, *Salvia macrosyphon* and *salvia rhytidea* were extracted by conventional methods [18-22]. Table 1 shows the species of Lamiaceae family extracted by conventional methods. *Mentha longifolia* and *Stachys lavandulifolia* were extracted by SFME method, *Mentha piperita* was extracted by MADD and MAHD and *Ocimum basilicum* was extracted by MAE and SFME [23-28]. Table 2 shows the species of Lamiaceae family extracted by different microwave extraction methods. For all these plants the aerial parts were utilized for essential oil extraction in both HD and microwave extraction methods, but in case of *Salvia macrosyphon* and *salvia rhytidea* the seeds also were used for essential oil extraction.

Nepeta is the largest genera of the Lamiaceae family, and extensively used in folk medicine because of their antispasmodic, diuretic, antiseptic, antitussive and antiasthmatic activities. Most *Nepeta* species are rich in essential oils [29]. In Afghanistan, genus *Nepeta*

comprises of 49 species in which 17 of them are endemic.

Salvia is another large genus in Lamiaceae family comprise of many species which have been widely used in traditional medicine therapy. A large amount of genus *salvia* is economically important and used as spices and flavouring agent [30]. The therapeutic properties of *Salvia* are related to their essential oils. The species of *Salvia* bear the attributes of antioxidant, antimicrobial, antifungal, and aromatic [31]. In Afghanistan 25 species of this genus exist over them five species are endemic and have not been studied yet. Table 3 shows the endemic species of Afghanistan belonging to genus *Nepeta*, *Salvia*, *Mentha* and *Thymus*.

3.2 Apiaceae (Umbelliferae)

Apiaceae is a large botanical family which composed of 300-455 genera and more than 3,000 species. The Apiaceae family is usually aromatic plant. Therefore, various Umbelliferae genera are used in industries because of their aromatic and medical virtues [32, 33].

In Afghanistan, 214 species of Apiaceae family are found in which 56 of them are endemic [1]. Totally, 15 plant of Umbelliferae family are introduced as medical plant [2]. Thirteen of these plants contain essential oils. *Conium maculatum*, *Daucus carota*, *Dorema ammoniacum*, *Ferula foetida*, *Prangos pabularia* are extracted by conventional methods [34-38]. Different microwave extractions were applied for essential oil extraction of *Anethum graveolens*, *Opium graveolens*, *Carum carvi*, *Carum copticum*, *Coriandrum sativum*, *Cuminum cyminum*, *Foeniculum vulgare* and *Pimpinella anisum* [39-49]. Tables 1 and 2 show the species of Apiaceae family extracted by conventional and microwave extraction methods. Almost the seeds of these plants were treated for extracting essential oil expect for *C. Maculatum*, *O. graveolens* and *A. graveolens* whose leaves and aerial parts were used for oil extraction.

Table 1 Afghanistan medicinal plants applied to conventional extraction methods.

Local name in Afghanistan	Botanical name	Family name	Parts used for study	Bioactivity	Extraction method
Bozbash	<i>Nepeta laevigata</i> (Don) Hand. Mzt.	Lamiaceae	Leaves	Fever, Sore throat,	SD [18]
Surkh Sawij	<i>Salvia hydrangea</i> L.	Lamiaceae	Aerial part	Carminative, Spasmolytic, anti-inflammation	HD [19]
Kanawcha	<i>Salvia macrosyphon</i> Boiss.	Lamiaceae	Aerial part	Emollient, anti-tussive	HD [20] [21]
Malangan	<i>Salvia rhytidea</i> Benth.	Lamiaceae	Seeds Flowers Leaves	Expectorant, anti-tussive, emollient	HD [22]
Margig	<i>Conium maculatum</i> L.	Umbeliferae (Apiaceae)	Leaves Flowers	Toxic	HD [34]
Zardak	<i>Daucus carota</i> L.	Umbeliferae (Apiaceae)	Seeds	Column analgesic, Diuretic, Stomachic	SD [35]
Ganda Firoza	<i>Dorema ammoniacum</i> Don.	Umbeliferae (Apiaceae)	Fruits	Antimicrobial activity	HD [36]
Heng	<i>Ferula foetida</i> (bunge) Regel.	Umbeliferae (Apiaceae)	Gum-Resin	Spasmolytic, anthelmintic	HD [37]
Burboo	<i>Prangos pabularia</i> Lindl.	Umbeliferae (Apiaceae)	Leaves Fruits Umbel	Antiseptic, diuretic, aphrodisiac, digestive disorders, scars, bleeding	HD [38]
Boemadaran	<i>Achillea santolina</i> L.	Compositae	Aerial part	Antimicrobial, anti-inflammatory, antigastritis	HD [50]
Aqha Anqara	<i>Anacyclus pyrethrum</i> L.	Compositae	Root Aerial part	Antimicrobial activity, analgesic, antigastritis, anthelmintic	HD [51]
Mastar	<i>Artemisia alba</i> L.	Compositae	Aerial part	Antipyretic, diuretic, anthelmintic, anti-diabetic, leishmanicidal, antibacterial, antifungal	HD [52] [53]
Bahman Safid	<i>Centaurea behen</i> L.	Compositae	Aerial part	Cytotoxic, antibacterial, anti-inflammatory, hypotensive	HD [54]
Pirtaran	<i>Chrysanthemum parthenium</i> Pers.	Compositae	Flowers	Anti-migraine, anti-arthritis, anti-psoriasis, antibacterial, antioxidant, insecticide	HD [55]
Kasni	<i>Cichorium intybus</i>	Compositae	Aerial part	Anti-hypototic, anti-diabetic, anti-malaria antipyretic	HD [56]
Zanjibir Shahi	<i>Inula helenium</i> L.	Compositae	Roots	Anti-bacterial, cerebral analgesic	HD [57]
Kahoo	<i>Lactuca sativa</i> L.	Compositae	Seeds Leave	Anti-microbial, antifungal, anti-bacterial, sedative, branchopulmonary infection	HD [58] [59]
Zrad Sarak	<i>Matricaria disciformis</i> DC. <i>Tripleurospermum disciformis</i> (C. A. Mey.) Sch. Bip.	Compositae	Aerial part	Gastrotonic, carminative, tranquilizer, antifungal, hair tonic, antihemorrhage	HD [60]

(table 1 continued)

Local name in Afghanistan	Botanical name	Family name	Parts used for study	Bioactivity	Extraction method
Parisiawashan	<i>Adiantum capillus-veneris</i> L.	Adiantaceae	Leaves	Expectorant, diuretic, laxative, anti-diarrheal	HD [61]
Kalpura	<i>Aerva javanica</i> (Burn. f.) Spreng.	Adiantaceae	Leaves Stem Seeds	Anti-gastritis, anti-diabetic, diuretic, sedative	Dry SD [62] HD [63]
Khenjak	<i>Pistacia khinjuk stocks</i> .	Anacardiaceae	Aerial part	Sedative, digestion disorders, tonic, toothache, astringent	HD [64]
Gule Gowzuban	<i>Echium amoenum</i> L.	Betulaceae	Flowers	Tonic, tranquilizer, diaphoretic, anti-pneumonia, cough suppressant	HD [65]
Korgiah	<i>Capparis spinosa</i> L.	Capparaceae	Aerial part	Analgesic, anthelmintic, expectorant, diuretic	HD [66]
Chambli	<i>Lonicera caprifolium</i> DC.	Caprifoliaceae	Flowers	Anti-bacterial, antiviral, antioxidant	HD [67]
Awri (Khardal)	<i>Brassica hirta moench</i> .	Cruciferae	Seeds	Bronchopulmonary infection, menstruation disorders, anti-neuralgia, anti-pneumonia	SD [68]
Tarboz Abujahel	<i>Citrullus colocynthis schrad</i> .	Cucurbitaceae	Seeds	Laxative, anti-cathartic, antioxidant, anti-hypersensitive, anti-diabetic, immunostimulant, anti-bacterial	HD [69]
Abhal	<i>Juniperus sabina</i> L.	Cupressaceae	Fruit Leaves	Anti-neoplastic, abortive, antibacterial, antifungal	SD [70]
Amla	<i>Emblica officinalis gaertn</i> .	Euphorbiaceae	Fruits Seeds	Anti-bacterial, diuretic, laxative, hair tonic, anti-insomnia, anti-hemorrhage	SD [71]
Bed Anjir	<i>Ricinus communis</i> L.	Euphorbiaceae	Aerial part	Anti-microbial, anti-inflammation, anti-diabetic, liver disorders, laxative	HD [72]
Shahtara	<i>Fumaria parviflora</i> Lam.	Fumariaceae	Aerial part	Anti-histaminic, insecticidal	HD [73]
Shirinboya	<i>Glycyrrhiza glabra</i> L.	Leguminosae	Leaves	Anti-inflammatory, anti-fungal, anti-cancer, antioxidant, anti-bacterial	HD [74]
Shanbalilia (Hulba)	<i>Trigonella Foenum Graecum</i> L.	Leguminosae	Aerial part Seeds	Anti-diabetic, anti-fever, antioxidant, anti-inflammation, anti-microbial	HD [75] SE [76]
Anjir	<i>Ficus carica</i> L.	Moraceae	Leaves	Anti-hemoroidal, anti-anemic, stimulant	HD [77]
Bartang	<i>Plantago lanceolata</i> L.	Plantagianaceae	Fruit Leaves	Emollient, antitussive	HD [78] SE
Anjabar	<i>Polygonum bistorta Gaecke</i> .	Polygonaceae	Flowers	Antioxidant, refreshing, stimulant	HD [79]
Anar	<i>Punica granatum</i> L.	Portulacaceae	Seeds Flowers	Anti-diabetic, astringent, anti-diarrhea	CP/HD [80] [81]
Gulab	<i>Rosa centifolia</i> L.	Rosacea	Flowers	Mild Anti-viral and bactericidal, cooling, relaxing, toning, anti-asthma	SE [82]
Fuwa	<i>Rubia tinctorum</i>	Rubiaceae	Aerial part	Amenorrhea, dropsy, jaundice	HD [83]

(table 1 continued)

Local name in Afghanistan	Botanical name	Family name	Parts used for study	Bioactivity	Extraction method
Sadab	<i>Ruta graveolens</i> L.	Rutaceae	Aerial part	For digestive disorders	HD [84]
Rita	<i>Sapindus trifoliatus</i> L.	Santalaceae	Seed	Detergent, surfactant, emulsionant	HD [85]
Murche Surkh	<i>Capsicum annuum</i>	Solanaceae	Fruits	Flavoring agent, Food dye	SD [86]
Datura	<i>Datura stramonium</i>	Solanaceae	Aerial part	Anti-bacterial, antioxidant, spasmolytic, anti-asthmatic, aphrodisiac	SD [87]
Sag Angorak	<i>Solanum nigrum</i> L.	Solanaceae	Leaves Aerial parts	Anti-ulcer, analgesic, sedative, anti-tuberculoses, anti-tumor, antioxidant, anti-inflammatory	HD [88]
Panirband	<i>Withania coagulans</i> Dunal.	Solanaceae	Fruits	Emetic, stomachic, anti-diabetic	SD [89]
Susan bekh (Benafsha)	<i>Viola odorata</i> L.	Violaceae	Aerial part	Anti-inflammatory, expectorant, diuretic, antioxidant, as perfume	SD [90]
Espand	<i>Peganum harmala</i> L.	Zygophyllaceae	Leaves Seeds	Erosive, hypnotic, antispasmodic, anodyne, emetic	HD [91] [92]
Benafsha Tokhom(Lale Sarnegon)	<i>Fritillaria imperialis</i> L.	Liliaceae	Aerial part	Cardiac stimulant	HS [93]
Koknar	<i>Papaver somniferum</i> L. Var. Album L.	Papaveraceae	Seeds	As adjuvant for medical diagnostics, as a carrier for cancerostatics in the treatment of hepatocellular carcinoma and cyclosporine A.	SPME [94]
Bang Dana	<i>Cannabis Sativa</i> L.	Cannabaceae	Bud	Narcotic	SD [95]
HD	Hydro-distillation				
SD	Steam distillation				
SE	Steam extraction				
SPME	Solid phase micro extraction				
CP/HD	Cold Press/Hydro-distillation				

Table 2 Afghanistan medicinal plants applied to microwave extraction methods.

Local name in Afghanistan	Botanical name	Family name	Part used for study	Bioactivity	Extraction methods
Raihan (Nazebo)	<i>Ocimum basilicum</i> L.	Lamiaceae	Aerial part	Antimicrobial, antioxidant	MAE [12] SFME [15]
Pudina	<i>Mentha longifolia</i> (L.) Huds.	Lamiaceae	Aerial part	Anti-inflammation, carminative, antiemetic, diaphoretic, antispasmodic, analgesic, anticatharral, stimulant	SFME [23] [24]
Nana	<i>Mentha piperita</i> L.	Lamiaceae	Leaves	Antiseptic, smoothing, antispasmodic, tonic, vasodilator	MADD [25] OAHD&MAHD [26]
Zarafshan	<i>Stachys lavandulifolia</i> vahl.	Lamiaceae	Aerial part	Anxiolytic, sedative	MAHD [27] [28]
Shebet	<i>Anethum graveolens</i> L.	Umbeliferae (Apiaceae)	Leaves Seed	Anti-fungal, diuretic	MAE [39]
Ajmood (Karafs)	<i>Opium graveolens</i> L.	Umbeliferae (Apiaceae)	Aerial parts	Carminative, stomachic, diuretic, emmenagogue, anti-rheumatism	MAE [40]
Zire Siah	<i>Carum carvi</i> L.	Umbeliferae (Apiaceae)	Seed	Perfumes, fragrances, spices, digestive disorders	MAE [41] MDG [42] MAWD [43]
Badian Sabez	<i>Pimpinella anisum</i> L. <i>Anisum vulgare</i>	Umbeliferae (Apiaceae)	Seeds	Carminative, antispasmodic, antiseptic, expectorant, stomachic, diuretic, diaphoretic, stimulant	MAWD [43]
Jawani	<i>Carum copticum</i> benth.	Umbeliferae (Apiaceae)	Seed	Antispasmodic, stimulant, tonic, carminative, antidiarrheal	SFME [44]
Gashniz	<i>Coriandrum sativum</i> L.	Umbeliferae (Apiaceae)	Leaves Seeds	Analgesic, antispasmodic, febrifuge, carminative, digestive	MAHD [45] MAE [46]
Zire Asel	<i>Cuminum cyminum</i> L.	Umbeliferae (Apiaceae)	Seeds	Spices, column analgesic, antioxidant	MAWD [43] SFME [44] [47] ISFME [46] MAE [48]
Badian Raziana	<i>Foeniculum vulgare</i> gaertn.	Umbeliferae (Apiaceae)	Fruits	Carminative, antioxidant, antibacterial, antifungal, mosquito repellent	MAWD [43] MAE [48] IMAE [49]
Afsantin	<i>Artemisia absinthium</i> L.	Compositae	Aerial part	Antifungal, antimicrobial, carminative, digestive disorders	SFME [96]
Hublqurtom (Masor)	<i>Carthamus tinctorius</i> L. (Safflower)	Compositae	Flowers	Analgesic, antispasmodic, emollient, anti-arthritis, cardiovascular, hypotensive	MD/SPME [97] [98]

(table 2 continued)

Local name in Afghanistan	Botanical name	Family name	Part used for study	Bioactivity	Extraction methods
Gole Babuna	<i>Matricaria chamomilla</i> L.	Compositae	Flowers	Antibacterial, antifungal, antiviral, antiparasitic, spasmolytic, antioxidant	MAE [99] [100]
Pista	<i>Pistacia vera</i> L.	Anacardiaceae	Fruit Leaves Gum	Antidiarrheal, sedative, alimentary	MAHD [101]
Archa dana (Sarwe Kohi)	<i>Juniperus excelsa</i> bieb.	Cupressaceae	Aerial part	Menstrual analgesia anti-tussive, anti-bronchitis, anti-tuberculosis	SFME [102] [103]
Zafaran	<i>Crocus sativus</i>	Iridaceae	Stigma Corm	Antispasmodic, expectorant, aphrodisiac	MAE [104]
Henna (Khina)	<i>Lawsonia inermis</i> roxb.	Lythraceae	Leaves	Cosmetics, antioxidant	MAHT [105] [106]
Siah dana	<i>Nigella sativa</i> L.	Ranunculaceae	Seeds	Anti-asthmatic, anti-tumor, antiviral, antibacterial, anti-inflammatory, anti-malarial, antihypertensive, anti-diabetic, anti-atherosclerotic, gastroprotective, antioxidant, anti-cholesterol	MSD [107]
Samaruq	<i>Rhus coriaria</i> L.	Therebintaceae	Fruits Leaves Flowers	Antiseptic, food flavoring agent, antioxidant, antimicrobial	MAE [108]
Panj Angusht	<i>Vitex negundo</i> L.	Verbenaceae	Leaves	Anthelmintic, antibacterial	SFME [109] MAHD

MAHD	Microwave Assisted Hydro-distillation
MSD	Microwave Steam Distillation
MASE	Microwave Assisted Steam Extraction
MAHT	Microwave Assisted Hydro-thermal Extraction
MD/SPME	Microwave Distillation/Solid Phase Micro extraction
OAHD	Ohmic Assisted Hydro-distillation
ISFME	Improved Solvent Free Microwave Extraction
MAWD	Microwave Assisted Water Distillation
SFME	Solvent Free Microwave Extraction

Table 3 The endemic species of Afghanistan plant in the genera of three botanical families comprise essential oil.

No.	Compositae		Apiaceae		Lamiaceae	
	Species	Genus	Species	Genus	Species	Genus
1	<i>A. andersiana</i>	Artemisia	<i>F. afghanica</i>	Ferula	<i>N. amicomum</i>	Nepeta
2	<i>A. bicolor</i>	Artemisia	<i>F. costata</i>	Ferula	<i>N. barfakensis</i>	Nepeta
3	<i>A. dumosa</i>	Artemisia	<i>F. dictyocarpa</i>	Ferula	<i>N. bellevii</i>	Nepeta
4	<i>A. ghazniensis</i>	Artemisia	<i>F. ghorana</i>	Ferula	<i>N. freitagii</i>	Nepeta
5	<i>A. ghoratensis</i>	Artemisia	<i>F. glabra</i>	Ferula	<i>N. juncea</i>	Nepeta
6	<i>A. Kandaharensis</i>	Artemisia	<i>F. hedgeana</i>	Ferula	<i>N. nawarica</i>	Nepeta
7	<i>C. codringtonii</i>	Centaurea	<i>F. heratensis</i>	Ferula	<i>N. paktiana</i>	Nepeta
8	<i>C. heratensis</i>	Centaurea	<i>F. kandahrica</i>	Ferula	<i>N. persica</i>	Nepeta
9	<i>I. sericeo-villosa</i>	Inula	<i>F. myrioloba</i>	Ferula	<i>N. podlechii</i>	Nepeta
10			<i>F. nuristanica</i>	Ferula	<i>N. polyodonta</i>	Nepeta
11			<i>F. pachycaulos</i>	Ferula	<i>N. rechingeri</i>	Nepeta
12			<i>F. rechingeri</i>	Ferula	<i>N. subincisa</i>	Nepeta
13			<i>F. stenoloba</i>	Ferula	<i>N. uberrima</i>	Nepeta
14			<i>F. trachelocarpa</i>	Ferula	<i>S. ariana</i>	Salvia
15			<i>F. trachyphylla</i>	Ferula	<i>S. maymonica</i>	Salvia
16			<i>F. xanthocarpa</i>	Ferula	<i>S. pterocalyx</i>	Salvia
17					<i>S. rechingeri</i>	Salvia
18					<i>S. tetradonta</i>	Salvia
19					<i>M. longifolia</i> (<i>austroafghanica</i>)	Mentha
20					<i>M. royleana</i> (<i>afghanica</i>)	Mentha
21					<i>T. koeieanus</i>	Thymus

The genus *Ferula*—the old-world plant, belongs to Apiaceae family and has some 130 species distributed throughout the Mediterranean area and Central Asia. The plants of this genus are good source of biologically active compounds such as derivatives and sulfur containing compounds [33]. These plants are frequently used as spices and in the provision of local drugs. Some species are used in folk medicine for the treatment of skin infections and hysteria [110]. The Afghanistan flora comprises of 31 species of *Ferula*, of which 16 of them are endemic. Table 3 shows the endemic species of *Ferula* genus in Afghanistan [1].

3.3. Compositae (*Asteraceae*)

Compositae botanical family contains 9,000 genera and about 20,000 species. Over 180 species are used for medical purposes [111].

In Afghanistan, 705 species of Compositae family are found in which 192 of them are endemic [1]. Shafique et al. [2] introduced seventeen species of

Compositae family as medical plants of Afghanistan. Among them, fifteen species are containing essential oil. *Achillea santolina*, *Anacyclus pyrethrum*, *Artemisia alba*, *Centaurea behen*, *Chrysanthemum parthenium*, *Cichorium intybus*, *Inula helenium*, *Lactuca sativa* and *Matricaria disiformis* are extracted by HD [50-60]. *Artemisia absinthium*, *Carthamus tinctorins* and *Matricaria chamomile* are extracted by different microwave extraction methods [96-100]. Essential oil in the Compositae family is distributed in all parts of the plants. In *A. santolina*, *C. behan*, *Artemisia* genus, *C. intybus*, and *M. disiformis* the essential oil was extracted from aerial part while for *A. pyrethrum*, *I. helenium*, *L. sativa* seeds and for *C. parthenium*, *C. tinctorins* and *M. chamomile* flowers were used for extraction. Tables 1 and 2 show conventional and microwave extraction of plants belong to Compositae family.

Genus *Artemisia* is one of the most abundant plants in Compositae family in the world and has special

interest because of botanical and pharmaceutical properties. Terpenoids are one of the major constituents of this genus which makes it the most momentous source of biological compounds [112]. In Afghanistan, the genus *Artemisia* consists of 54 species, six of which are endemic [1]. Table 3 shows the endemic species of *Artemisia* genus in Afghanistan.

4. Essential Oil in Afghanistan

A number of small companies in Afghanistan produce essential oils from bitter orange blossom (Gule narinj) and rose which are used by perfume industries in France [113]. As well as, in Afghanistan, much amount of essential oil is extracted from medicinal plants by boiling water and making tea or by adding seeds or aerial parts of plants in food and cooking them in the kitchen in everyday life of normal people. In these methods less amount of essential oil is extracted. A conventional extractor is specially designed and is difficult to use in houses, but nowadays microwave ovens are available in many markets with reasonable prices and some people already use in their kitchen in cities [114].

5. Conclusions

The purpose of this review was to show the importance of Afghanistan medicinal plants and their utilization in domestic and industrial areas. Two methods of extraction: microwave and conventional extraction applied widely for extraction of essential oil from plant were reviewed in this article.

Species from Compositae, Lamiaceae and Apiaceae which contain much amount of essential oils are widely spread in Afghanistan and traditionally used by people as food, spices and medicines.

Extractions of essential oils from plants which are not endemic plants of Afghanistan have been done in other countries. As the yield and chemical composition of essential oils affected by growth place of a single species, study of species of botanical families from Afghanistan is highly recommended.

Also the native flora in these families still includes many plants for instance genus *Artemisia* which may contain essential oil, have not been studied yet and can result in new products for pharmaceuticals, perfumes and cosmetics industries.

Given the fact that microwave energy is increased the extracted amount of essential oil and the time of extraction decreased. Therefore, this study may encourage domestic people to use a microwave oven for extracting much amount of essential oil in their kitchen in very short time and take the advantage of both food and medicine from the plants which they use in their daily life as food.

Acknowledgments

The authors greatly acknowledge Professor Mohammad Usman Babory of Kabul University for his guidance in writing this paper. Also we would like to thank Mr. Rasool Hossaini for making available his bachelor monograph regarding traditional usage of medicinal plants by residents of Behsood District of Afghanistan, for us.

References

- [1] Breckel, S. W., Hedge, I. C., and Rafiqpoor, M. D. 2013. *Vascular plants of Afghanistan: an augmented checklist*. p. 25.
- [2] Shafique, Y., Jacques, F., Dominique, N., Guy, M., Francois, M., and Jean, M. P. 1987. "Repertory of Drugs and Medicinal Plants Used in Traditional Medicine of Afghanistan." *J. Ethnopharmacol* 20: 245-90.
- [3] Jeppesen, A. S., Soelberg, J., and Jager, A. K. 2012. "Antibacterial and COX-1 Inhibitory Effect of Medicinal Plants from the Pamir Mountains, Afghanistan." *Plants* 1: 74-81.
- [4] Kassam, A. K., Karamkhudoeva, M., Ruelle, M., and Baumflek, M. 2010. "Medical Plant Use and Health Sovereignty: Findings from Tajik and Afghan Pamir." *Hum. Ecol.* 38: 817-29.
- [5] Edris, A. E. 2007. "Pharmaceutical and Therapeutical Potentials of Essential Oils and Their Individual Volatile Constituents." *Phytother. Res.* 21: 308-23.
- [6] Boelens, M. H. 1985. "Essential Oils and Aroma Chemicals from *Eucalyptus Globulus* Labill." *Perfumer & Flavorist* 9: 1-14.

- [7] Luque de Castro, M. D., Jimenez-Carmona, M. M., and Perez, V. F. 1999. "Towards More Rational Techniques for the Isolation of Valuable Essential Oils from Plants." *Trends Anal. Chem.* 18: 708-16.
- [8] Megha, J., Trisa, P., and Bhagat, S. 2014. "Variation in the Vitamin C and Essential Oil Content of *Ocimum Sanctum*, L, Growing in the Various Regions of Kerala and Maharashtra." *Inter. J. Scientific Res.* 3: 34-6.
- [9] Saoud, A. A., Yunus, R. M., Aziz, R. A., and Rahmat, A. R. 2005. "Study of Eucalyptus Essential Oil Acquired by Microwave Extraction." *Acta Hort* 679: 173-9.
- [10] Lucchesi, M. E., Chemat, F., and Smadja, J. 2004. "Solvent Free Microwave Extraction: An Innovation Tool for Rapid Extraction of Essential Oil from Aromatic Herbs and Spices." *J. Microw. Power Electromagn. Energy* 39: 135-9.
- [11] Okoh, O. O., Sadimenko, A. P., and Afolayan, A. J. 2010. "Comparative Evaluation of the Antibacterial Activities of the Essential Oils of *Rosmarinus officinalis* L. Obtained by Hydrodistillation and Solvent Free Microwave Extraction Methods." *J. Food Chem.* 120: 308-12.
- [12] Ugarte, G. A., Becerra, G. P., Morales, M. E., and Malo, A. L. 2013. "Microwave-Assisted Extraction of Essential Oils from Herbs." *J. Microw. Power Electromagn. Energy* 47: 63-72.
- [13] Chan, C. H., Yusoff, R., Ngoh, G. C., and Kung, F. W. L. 2011. "Review: Microwave-Assisted Extraction of Active Ingredients from Plants." *J. Chromatogr. A* 1218: 6213-25.
- [14] Vian, M. A., Fernandez, X., Visinoni, F., and Chemat, F. 2008. "Microwave Hydrodiffusion and Gravity: A New Technique for Extraction of Essential oils." *J. Chromatogr. A* 1190: 14-7.
- [15] Lucchesi, M. E., Chemat, F., and Smadja, J. 2004. "Solvent-Free Microwave Extraction of Essential Oil from Aromatic Herb: Comparison with Conventional Hydro-Distillation." *J. Chromatogr* 1043: 323-7.
- [16] Bernath, J. 2009. "Aromatic Plants." In *Cultivated plants, Primarily as Food Sources* 2: 329-52.
- [17] Zamfirachi, M. M., Burzo, I., Padurariu, C., Boz, I., Andro, R. A., Badea, L. M., et al. 2010. "Studies Regarding the Chemical Composition of Volatile Oils from Some Spontaneous and Cultivated Lamiaceae Species." *Biologie Vegetala* 43-9.
- [18] Hassan, T., Rather, M. A., Shawl, A. S., Bhat, K. A., Haroon, M., Bhat Dar, B. A. et al. 2011. "Chemical Composition of the Essential Oils of *Nepeta Laevigata* and *Nepeta Elliptica* from India." *Chem. Nat. Compd.* 47: 456-8.
- [19] Ghannadi, A., Shariat, S. H., and Moattar, F. 1999. "Volatile Constituents of the Flower of *Salvia Hydrangea* DC." *EX Benth. Daru* 7: 23-5.
- [20] Sefidkon, F., Mirza, M., and Javidtash, I. 2005. "Essential Oil Composition of *Salvia Macrosiphon Boiss* from Iran." *J. Essent. Oil Bear Plants* 8: 126-9.
- [21] Rowshan, V., and Bijeli, M. 2013. "Comparison of Chemical Constituents of Essential oil and Antioxidant Activity of *Salvia Macrosiphon Bioos*. (Wild and Cultivated Type) by FRAP and DPPH Assay." *Int. J. Agron. Plant Prod.* 4: 1197-203.
- [22] Sajjadi, S., and Ghannadi, A. 2005. "Essential Oil of the Persian Sage, *Salvia Rhytidea* Benth." *Acta Pharm* 55: 321-6.
- [23] Sharopov, F. S., Sulaimonova, V. A., and Setzer, W. N. 2012. "Essential Oil Composition of *Mentha Longifolia* from Wild Populations Growing in Tajikistan." *J. Med. Active Plants* 1: 76-84.
- [24] Okoh, O. O., and Afolayan, A. J. 2011. "The Effect of Hydrodistillation and Solvent Free Microwave Extraction Method on the Chemical Comparison and Toxicity of Essential Oils from the Leaves of *Mentha longifolia* L. subsp." *Capensis. Afr. J. Pharm. Pharmacol* 5: 2474-8.
- [25] Mircioaga, N., and Calinescu, I. 2011. "Extraction and Identification of Active Principles from *Mentha piperita* L." *Rev. Chim. (bucuresti)* 62: 1073-6.
- [26] Gavahian, M., Farahnaky, A., Farhoosh, R., Javidnia, K., and Shahidi, F. 2015. "Extraction of Essential Oils from *Mentha Piperita* Using Advanced Techniques: Microwave versus Ohmic Assisted Hydrodistillation." *Food Bioprod. Process* 4: 50-8.
- [27] Faizbaksh, A., Tehrani, M. S., and Rustaiyan, A. 2003. "Composition of the Essential Oil of *Stachys Lavandulifolia* Vahl from Iran." *J. Essent. Oil Res.* 15: 72-3.
- [28] Sadarmomtaz, A., Meshkatsadat, M. H., and Taherparwar, P. 2011. "Comparison of Volatile Composition of *Stachys Lavandulifolia* Vahl Obtained by MWHD and HD Techniques." *Dig. J. Nanomater Biostruct* 6 (3): 1343-8.
- [29] Asgarpanah, J., Sarabian, S., and Ziarati, P. 2013. "Essential Oil of *Nepeta* Genus (Lamiaceae) from Iran: A Review." *J. Essen. Oil Res.* 1-12.
- [30] Fu, Z., Wang, H., Hu, X., Sun, Z., and Han, C. 2013. "The Pharmacological Properties of Essential Oils." *J. Appl. Pharm. Sci.* 3: 122-7.
- [31] Taarit, M. B., Msaada, K., Hosni, K., and Marzouk, B. 2014. "GC Analysis of *Salvia* Seeds as Valuable Essential Oil Source." *J. Adv. Chem.* 1-6.
- [32] Tabanca, N., Ma, G., Pasco, D. S., Bedir, E., Kirimer, N., Baser, K. H. C., et al. 2007. "Effect of Essential Oils and Isolated Compounds from *Pimpinella* Species on NF-kB: A Target for Anti-inflammatory Therapy." *Phytother Res.* 21: 741-5.
- [33] Sahebkar, A., and Iranshahi, M. 2010. "Biological

- Activities of Essential Oils from the Genus *Ferula* (Apiaceae)." *Asian Biomed* 4: 835-47.
- [34] Radulovic, N., Zlatkovic, D., Zlatkovic, B., Djokovic, D., Stojanovic, G., and Palic, R. 2008. "Chemical Composition of Leaf and Flower Essential Oils of *Conium Maculatum* from Serbia." *Chem. Nat. Compd.* 44: 390-2.
- [35] Chizzola, R. 2010. "Composition of the Essential Oil from *Daucus carota* ssp. *Carota* Growing Wild in Vienna." *J. Essent. Oil Bear Plants* 13: 12-9.
- [36] Yousefzadi, M., Mirjalili, M. H., Alnajjar, N., Zeinali, A., and Parsa, M. 2011. "Composition and *in vitro* Antimicrobial Activity of the Essential Oil of *Dorema Ammoniacum* D. Don. Fruit from Iran." *J. Serb. Chem. Soc.* 76: 857-63.
- [37] Kavossi, G., and Rowshan, V. 2013. "Chemical Composition, Antioxidant and Antimicrobial Activities of Essential Oil Obtained from *Ferula Ass-Foetida* Oleo-Gum-Resin: Effect of Collection Time." *Food Chem.* 38: 2180-7.
- [38] Razavi, S. M. 2012. "Chemical and Allelopathic Analysis of Essential Oils of *Prangos pabularia* Lindl. from Iran." *Nat. Prod. Res.* 26: 2148-51.
- [39] Stan, M., Lung, I., Opris, O., and Soran, M. L. 2014. "High-Performance Thin-Layer Chromatography Quantification of Some Essential Oils from *Anethum Graveolens* Extracts." *J. Planar. Chromatogr.* 27: 33-7.
- [40] Moradalizadeh, M., Akhgar, M. R., and Zarandi, F. 2013. "Microwave-Assisted and Conventional Hydrodistillation of Essential Oils from *Opium graveolens* L." *Asian J. Chem.* 25: 79-81.
- [41] Chemat, S., Amar, H. A., Lagha, A., and Esveld, D. C. 2005. "Microwave-Assisted Extraction Kinetics of Terpens from Caraway Seeds." *Chem. Eng. Process* 44: 1320-6.
- [42] Farhat, A., Tixier, A. S., Visinoni, F., Romdhane, M., and Chemat, F. 2010. "A Surprising Method for Green Extraction of Essential Oil from Dry Spices: Microwave Dry-Diffusion and Gravity." *J. Chromatogr. A* 1217: 7345-50.
- [43] Rahman, A., Hilphy, S., Fekaiki, D. F., and Hussain, R. A. 2015. "Extraction of Essential Oils from Some Types of Umbeliferace Family Using Microwave-Assisted Water Distillation." *J. Biol. Agric. Healthc.* 5: 16-28.
- [44] Lucchesi, M. E., Chemat, F., and Smadja, J. 2004. "An Original Solvent Free Microwave Extraction of Essential Oils from Spices." *Flavour Fragr J.* 19: 134-8.
- [45] Sourmaghi, M. H. S., Kiaee, G., Golfakhrabadi, F., Jamalifar, H., and Khanavi, M. 2014. "Comparison of Essential Oil Composition and Antimicrobial Activity of *Coriandrum sativum* L. Extracted by Hydrodistillation and Microwave-Assisted Hydrodistillation." *J. Food Sci. Technol.* 52 (4): 2452-7.
- [46] Zhai, Y., Sun, S., Wang, Z., Cheng, J., Sun, Y., Wang, L., et al. 2009. "Microwave Extraction of Essential Oils from Dried Fruits of *Illicium Verum* Hook. F. and *Cuminum cyminum* L. Using Ionic Liquid as the Microwave Absorption Medium." *J. Sep. Sci.* 32: 3544-9.
- [47] Wang, Z., Ding, L., Li, T., Zhou, X., Wang, L., Zhang, H., et al. 2006. "Improved Solvent-Free Microwave Extraction of Essential Oil from Dried *Cuminum cyminum* L. and *Zanthoxylum Bungeanum* Maxim." *J. Chromatogr. A* 1102: 11-7.
- [48] Hammouda, F. M., Saleh, M. A., Azim, N. S., Shams, K. A., Ismail, S. I., Shahat, A. A., et al. 2013. "Evaluation of the Essential Oil of *Foeniculum Vulgare* Mill (Fennel) Fruits Extracted by Three Different Extraction Methods by GC/MS." *Afr. J. Tradit. Complement Altern. Med.* 11: 277-9.
- [49] Patel, J. B., Patel, B., Patel, R. K., and Patel, B. H. 2012. "Comparative Evaluation of Extraction Method for Extraction of Essential Oil from *Foeniculum Vulgare*." *J. Pharm. Sci. Bio. Scientific Res.* 2: 176-8.
- [50] Bader, A., Flamini, G., Cioni, P. L., and Morelli, I. 2003. "Essential Oil Composition of *Achillea santolina* L. and *Achillea Biebersteinii* Afan. Collected in Jordan." *Flavour Fragr. J.* 18: 36-8.
- [51] Selles, C., Dib, M. E. A., Djabou, N., Beddou, F., Muselli, A., Tabti, B., et al. 2013. "Antimicrobial Activity and Evolution of the Composition of Essential Oil from Algerian *Anacyclus pyrethrum* L. through the Vegetative Cycle." *Nat. Prod. Res.* 27: 2231-4.
- [52] Mohsen, H., and Ali, F. 2009. "Essential Oil Composition of *Artemisia Herb-Alba* from Southern Tunisia." *Molecules* 14: 1585-94.
- [53] Belhaddad, R., Amor, L., Barroso, J. G., Pdero, L. G., and Figueiredo, A. C. 2014. "Essential Oil from *Artemisia Herb-Alba* Asso Grown Wild in Algeria: Variability Assessment and Comparison with an Updated Literature Survey." *Arabian J. Chem.* 7: 243-51.
- [54] Esmaeili, A., and Khodadadi, E. 2012. "Volatile Compounds of Essential Oil *Centaurea behen* L. Grown in Iran." *J. Pharm. Sci.* 3: 8-11.
- [55] Shafaghat, A., Larijani, K., and Salimi, F. 2009. "Composition and Antibacterial Activity of the Essential Oil of *Chrysanthemum Parthenium* Flower from Iran." *J. Essent. Oil Bear Plants* 12: 706-13.
- [56] Haghi, G., Arshi, R., Ghazian, F., and Hosseini, H. 2012. "Chemical Composition of Essential oil of Aerial Parts of *Cichorium intybus* L. from Iran." *J. Essent Oil Bear Plants* 15: 2136.
- [57] Deriu, A., Zanetti, S., Sechi, L. A., Marongiu, B., Piras, A., Porcedda, S., et al. 2008. "Antimicrobial Activity of *Inula helenium* L. Essential Oil against Gram-Positive and Gram-Negative Bacteria and *Candida* Spp." *Int. J.*

- Antimicrob Agents* 31: 581-92.
- [58] Xu, F., and Wang, H. A. A. 2011. "Analysis of Essential Oil Extracted from *Lactuca Sativa* Seeds Growing in Xinjiang by GC-MS." 34: 1887-91.
- [59] Al Nomaani, R. S. S., Hossain, M. A., Weli, A. M., Al-Riyami, Q., and Al-Sabahi, J. N. 2013. "Chemical Composition of Essential Oils and *in vitro* Antioxidant Activity of Fresh and Dry Leaves Crude Extract of Medicinal Plant of *Lactuca sativa* L. Native to Sultanate of Oman." *Asian Pac. J. Trop. Biomed* 3: 353-7.
- [60] Javidnia, K., Miri, R., Soltani, M., and Khosravi, A. R. 2008. "Essential Oil Composition of *Tripleurospermum Disciforme* from Iran." *Chem. Nat. Compd.* 44: 800-1.
- [61] Khodaie, L., Esnaashari, S., and Moghaddam, S. B. 2015. "Essential Oil of Aerial Part of *Adiantum Capillus-Veneris*: Chemical Composition and Antioxidant Activity." *J. Nat. Pharm. Prod.* 10: 1-5.
- [62] Samejo, M. Q., Memon, S., Bhangar, M. I., and Khan, K. M. 2012. "Chemical Composition of the Essential Oil of *Aerva Javanica* Leaves and Stems." *Pak. J. Anal. Environ Chem.* 13: 48-52.
- [63] Samejo, M. Q., Memon, S., Bhangar, M. I., and Khan, K. M. 2013. "Comparison of Chemical Composition of *Aerva Javanica* Seed Essential Oils Obtained by Different Extraction Methods." *Pak. J. Pharm. Sci.* 26: 757-60.
- [64] Pirbalouti, A. G., and Aghaei, K. 2011. "Chemical Composition of Essential Oil of *Pistacia Khinjuk* Stocks Grown in Bakhtiari Zagross Mountains, Iran." *Electron J. Biol.* 7: 67-9.
- [65] Semnani, K. M., and Saeedi, M. 2005. "Essential Oil Composition of *Echium Amoenum* FISCH & C. A. Mey." *J. Essent Oil Bear Plants* 8: 61-4.
- [66] Muhaidat, R., Al-Qudah, M. A., Al-Shayeb, A., Jacob, J. H., Al-Jaber, H. I., Hussein, E., et al. 2013. "Chemical Profile and Antibacterial Activity of Crude Fractions and Essential Oils of *Capparis Ovata* Deaf, and *Capparis spions* L. (Capparaceae)." *Int. J. Integr. Biol.* 14: 39-47.
- [67] Ilies, D. C., Radulescu, V., and Dutu, L. 2014. "Volatile Constituents from the Flowers of Two Species of Honeysuckle (*Lonicera Japonica* and *Lonicera Caprifolium*)." *Farmacja* 62: 194-201.
- [68] Miyazawa, M., and Kawata, J. 2006. "Identification of the Main Aroma Compounds in Dried Seeds of *Brassica Hirta*." *J. Nat. Med.* 60: 89-92.
- [69] Doss, A., Vijayasanthi, M., Anand, S. P., Paruguna, V., and Venkataswamy, R. 2011. "Screening of Antimicrobial Activity of Essential Oil and Methanol Extracts of *Citrullus colocynthis* (L.) Schrad. South As." *J. Biol. Sci.* 1: 7-15.
- [70] Asili, J., Emami, S. A., Rahimizadeh, M., Fazly-Bazzaz, B. S., and Hassanzadeh, M. K. 2010. "Chemical and Antimicrobial Studies of *Juniperus sabina* L. and *Juniperus Foetidissima* Willd Essential Oil." *J. Essent. Oil. Bear Plants* 13: 25-36.
- [71] Saxena, R., and Patil, P. 2014. "*In vitro* Antibacterial Activity of *Emblica Officinalis* Essential Oil against *Staphylococcus Aureus*." *Int. J. Theor. Appl. Sci.* 6: 7-9.
- [72] Zarai, Z., Chobba, I. B., Mansour, R. B., Bekir, A., Gharsallah, N., and Kadri, A. 2012. "Essential Oil of the Leaves of *Ricinus communis* L: Invitro Cytotoxicity and Antimicrobial Properties." *Lipids Health Dis.* 11: 1-7.
- [73] Ghayoor, H. S., and Saeidi, K. 2015. "Antifeedant Activities of Essential Oils of *Satureja Hortensis* and *Fumaria Parviflora* against Indian Meal Moth *Plodia Intrapunctellar Hubner* (Lepidoptera: Pyralidae)." *Entomol Ornithol Herpetol* 4: 1-4.
- [74] Chouitah, O., Meddah, B., Aoues, A., and Sonnet, P. 2011. "Chemical Composition and Antibacterial Activities of Essential Oil from *Glycyrrhiza Glabra* Leaves." *J. Essent Oil Bear Plants* 14: 284-8.
- [75] Sohrevardi, N., and Sohrevardi, F. 2012. "Essential Oil Composition and Antioxidant Activity of *Trigonella foenum graecum* L. Plant." *Int. J. Agri. Crop. Sci.* 4: 793-7.
- [76] Muman, E. M., Ibrahim, S. J., and Al-Yassin, H. D. 2014. "Study of Chemical Composition and the Biological Active Compounds of *Nigella Sativa* and *Trigonella foenum-graecum* L. Seed." *J. Appl. Chem.* 43-5.
- [77] Ayoub, N., Sigab, A. N., Mostafa, N., and Schultze, W. 2010. "Volatile Constituents of Leaves of *Ficus carica* Linn. Grown in Egypt." *J. Essent Oil Bear Plants* 13: 316-21.
- [78] Fons, F., Rapior, S., Gargadennec, A., Andary, C., and Bessiere, J. M. 1998. "Volatile Components of *Plantago lanceolata* (Plantaginaceae)." *Acta Bot Gallica* 145: 265-9.
- [79] Iskender, N. Y., Gulec, C. A., Yucel, M., Sinek, K., and Yayli, N. 2011. "Analysis of the Essential Oil from the Flower of *Polygonum bistorta* L. subsp. *Carneum* (Koch)." *Asian J. Chem.* 23: 1940-2.
- [80] Bakkour, Y., Makhoul, S., El-Nakat, H., and El-Omar, F. 2011. "Chemical Analysis of the Essential Oils from *Punica granatum*, *Vitis vinifera* and *Cucurbita maxima* Seeds Growing in Lebanon by GC/MS." *J. Nat. Prod.* 4: 71-4.
- [81] Mekni, M., Flamini, G., Garrab, M., Hmida, R. B., Cheraief, I., Mastouri, M., et al. 2013. "Aroma Volatile Components, Fatty Acids and Antibacterial Activity of Four Tunisian *Funica granatum* L., Flower Cultivars." *Int. Crops Prod.* 48: 111-7.
- [82] Shabbir, M. K., Nadeem, R., and Mukhtar, H. 2009. "Physico-Chemical Analysis and Determination of Various Chemical Constituents of Essential Oil in *Rosa Centifolia*." *Park. J. Bot.* 41: 615-20.

- [83] Navaei, M. N., Mirza, M., and Dini, M. 2006. "Chemical Composition of the Essential Oils of *Rubia tinctorum* L. Aerial Parts from Iran." *Flavour Fragr J.* 21: 519-20.
- [84] Soleimani, M., Azar, P. A., Tehrani, M. S., and Rustaiyan, A. 2009. "Volatile Composition of *Ruta graveolens* L. of North of Iran." *World Appl. Sci. J.* 7: 124-6.
- [85] Mahmood, M. A. A., Asif, I. M., Alam, M. S., Islam, M. S., Akhter, S., Hossain, F., et al. 2013. "Extraction and Characterization of Oils from *Sapindus trifoliatus* Linn Seed of Different Origin of Bangladesh." *Merit Res. J. Environ. Sci. Toxicol.* 1: 99-104.
- [86] Kevresan, Z. S., Hrabovski, N. C., Kuhajda, K. N., Dukic, N. M., and Sakac, M. B. 2009. "Essential Oil Composition of Fresh and Dried Pepper Fruits (*Capsicum annum* L.)." *Food Feed Res.* 36: 29-34.
- [87] You, L. X., and Wang, S. J. 2011. "Chemical Composition and Allelopathic Potential of the Essential Oil from *Datura stramonium* L." *Adv. Mater. Res.* 233: 2472-5.
- [88] Ogundajo, A. L., Oldosu, I. A., Ogunwande, I. A., Flamini, G., and Owolabi, M. S. 2013. "Study on the Volatile Constituents of *Solanum nigrum* Var. *Virginicum* L. from Nigeria." *Asian J. Plant Sci. Res.* 3: 94-8.
- [89] Bakhtawar, S., Mughal, T., and Naeem, I. 2010. "Chemical Composition of the Essential Oil of *Withania Coagulans*." *Asian J. Chem.* 22: 122-6.
- [90] Akhbari, M., Batooli, H., and Kashi, F. J. 2012. "Composition of Essential Oil and Biological Activity of Extracts of *Viola odorata* L. from Central Iran." *Nat. Prod. Res.* 26: 802-9.
- [91] Faridi, P., Ghasemi, Y., and Mohagheghzadeh, A. 2013. "Chemical Composition of *Peganum Harmala* Smoke and Volatile Oil." *J. Essent. Oil Bear Plants* 16: 850-4.
- [92] Abdellah, K., Zakaria, B., Nawal, B., Aminata, O. K., and Didi, O. M. 2013. "Biological Activity of Essential Oils Leaves from One Sahara Plant: *Peganum harmala* L. (Zygophyllaceae) on the Desert Locust." *Int. J. Curr. Microbiol. App. Sci.* 2: 389-95.
- [93] Helsper, J. P., Bucking, M., Muresan, S., Blaas, J., and Wietsma, W. A. 2006. "Identification of the volatile Component(s) Causing the Characteristic Foxy Odor in Various Cultivars of *Fritillaria imperialis* L. (Liliaceae)." *J. Agric. Food Chem.* 54: 5087-91.
- [94] Krist, S., Stuebinger, G., Unterweger, H., Bandion, F., and Buchbauer, G. 2005. "Analysis of Volatile Compounds and Triglycerides of Seed Oils Extracted from Different Poppy Varieties (*Papaver somniferum* L.)." *J. Agric. Food Chem.* 53: 8310-6.
- [95] Ross, S. A., and Elsohly, M. A. 1996. "The Volatile Oil Composition of Fresh and Air-Dried Buds of *Cannabis Sativa*." *J. Nat. Prod.* 59: 49-51.
- [96] Tehrani, M. S., Azar, P. A., Hosain, S. W., Khalilzadeh, M. A., and Zanousi, M. B. 2012. "Composition of Essential Oil of *Artemisia Absinthium* by Three Different Extraction Methods: Hydrodistillation, Solvent-Free Microwave Extraction & Headspace Solid-Phase Microextraction." *Aisan J. Chem.* 24: 5371-6.
- [97] Ziarati, P., Asgarpana, J., and Kianifard, M. 2012. "The Essential Oils Composition of *Carthamus tinctorius* L. Flowers Growing in Iran." *Afr. J. Biotechnol.* 11: 12921-4.
- [98] Yu, Y., Yang, B., Zhou, T., Zhang, H., Shao, L., and Duan, G. 2007. "Rapid Determination of Volatile Constituents in Safflower by Microwave Distillation and Simultaneous Solid-Phase Microextraction Coupled with Gas Chromatography-Mass Spectroscopy." *Ann. Chem.* 97: 1075-84.
- [99] Kazemi, M. 2014. "Chemical Composition and Antimicrobial Activity of Essential Oil of *Matricaria Chamomilla*." *Bull. Env. Pharmacol. Life. Sci.* 3: 148-53.
- [100] Gomez, N. E., and Witte, L. 2001. "A Simple Method to Extract Essential Oils from Tissue Samples by Using Microwave Radiation." *J. Chem. Ecol.* 27: 2351-9.
- [101] Moghaddam, H. H., Mohammadhosseine, M., and Salar, M. 2014. "Chemical Composition of the Essential Oils from the Huls of *Pistacia vera* L. by Using Magnetic Nanoparticle-Assisted Microwave (MW) Distillation: Comparison with Routine MW and Conventional Hydrodistillation." *Anal. Methods* 6: 2572-9.
- [102] Unlu, M., Unlu, G. V., Vural, N., Donmez, E., and Cakmak, O. 2008. "Composition and Antimicrobial Activity of *Juniperus Excelsa* Essential Oil." *Chem. Nat. Compd.* 44: 129-31.
- [103] Hussain, J., Rehman, N. U., Al-Harrasi, A., Ali, L., Khan, A. L., and Albroumi, M. A. 2013. "Essential Oil Composition and Nutrient Analysis of Selected Medicinal Plants in Sultanate Oman." *Asian Pac. J. Trop. Dis.* 3: 421-8.
- [104] Masuda, A., Mori, K., Oda, Y., and Miyazawa, M. 2010. "Volatile Oil Compounds from Corms and Flowers of *Crocus vernus* L. Hill and Corms of *C. sativus* L." *Libyan Agric. Res. Cen. J. Int.* 1: 244-9.
- [105] Oyedeji, A. O., and Ekundayo, O. 2005. "Essential Oil Composition of *Lawsonia inermis* L. Leaves from Nigeria." *J. Essent. Oil Res.* 17: 403-4.
- [106] Zohourian, T. H., Quitain, A. T., and Sasaki, M. 2011. "Polyphenolic Contents and Antioxidant Activities of *Lawsonia Inermis* Leaf Extracts Obtained by Microwave-Assisted Hydrothermal Method." *J. Microw. Power Electromagn Energy* 45: 193-204.
- [107] Akloul, R., Ali, F. B., Zerrouki, M., and Eppe, G. 2014. "Composition and Biological Activities of the Essential Oil of *Nigella Sativa* Seeds Isolated by Accelerated Microwave Steam Distillation with Cryogenic Grinding." *Am. J. Essent. Oil Nat. Prod.* 1: 23-33.

**Extraction of Essential Oils from Afghanistan Medicinal Plants
Using Microwave and Conventional Methods**

- [108] Gharaei, A., Khajeh, M., Ghaffari, M., and Choopani, A. 2013. "Iranian Rhus Coriaria (Sumac) Essential Oils Extraction." *J. Essent. Oil Bear Plants* 16: 270-3.
- [109] Azar, P. A., Torabbeigi, M., and Tehrani, M. S. 2011. "The Investigation of Essential Oil of Vitex Pseudo-Negund with Different Analytical Methods: Hydrodistillation, SFME, HS-SPME and MA-HS-SPME." *J. Essent. Oil Bear Plants* 14: 755-60.
- [110] Zellagui, A., Gherraf, N., and Rhouati, S. 2012. "Chemical Composition and Antibacterial Activity of the Essential Oils of Ferula Vesceritensis Coss et Dur. Leaves, Endemic in Algeria." *Org. Med. Chem. Lett.* 2: 1-4.
- [111] Mollasalehi, S., Kashefi, B., and Moghaddam, H. H. 2013. "Comparison of Microwave-Assisted Hydrodistillation Methods for Extraction of Essential Oil from Achillea Millefolium." *J. Chem. Health Risk* 3: 39-46.
- [112] Pirbalouti, A. G., Craker, L., and Akbarzadeh, M. 2013. "Essential Oil Composition, Antibacterial and Antioxidant Activities of Various Populations of Artemisia Chamaemelifolia at Two Phenological Stages." *Rev. Bras. Farmacogn.* 23: 861-9.
- [113] "Essential Oils and Natural Perfumes: A Rural Development Opportunity in Afghanistan-Gulestan Ariana Company." In *Proceedings of the Enabling Environment Conference 2007*, pp. 129-34.
- [114] <https://afghanistan.jadopado.com/category/06064/microwave-ovens/>.